

NOAA FISHERIES

Northwest Fisheries Science Center

Effects of vessels on southern resident killer whale behavior

Dawn P. Noren
Conservation Biology Division
Northwest Fisheries Science Center


West Coast Marine Mammal Program Review July 2015

Vessel Presence



Studies that addressed management needs relevant to proposed vessel regulations


Voluntary Guidelines





Land-Based Theodolite SRKW Behavioral Study

David Bain (2003-2005)





- Scan sample (Lusseau et al. 2009)
 - ↓ Foraging and ↑ travel when boats are within 100 m and from 100 - 400 m of whales
- Focal follow (Williams et al. 2009)
 - Changes in respiration rate, swim speed, path directedness, and occurrence of surface active behaviors (SABs) when boats are within 0 to ≥400 m
 - ↑ Performance of SABs in response to close approaches by vessels and ↑ vessel presence



Boat-Based Focal Follow SRKW Behavioral Study

Dawn Noren (2004-2006)






- Spatial and temporal relationships between close approaches by vessels and surface active behaviors (SABs, Noren et al. 2009)
 - SAB bouts ↑ when vessel distance to whale ↓
 - Significantly higher proportion of SAB bouts occur when vessels are within 0 to 150 m of whales
 - SAB bouts ↑ near the time of the closest approach by a vessel

Boat-Based Scan Sample SRKW Behavioral Study

Dawn Noren (2006)









- Some SRKW activity states predominantly occur in distinct locations
 - Rest 2 locations
 - Forage Primarily southwest of San Juan Island

Persistent Activity State Hot Spots

2006 (Noren and Hauser in review)



1976-83 (Heimlich-Boran 1988)



Forage: Southwest of San Juan Island

Rest: Northwest of San Juan Island




Research Efforts Resulted in Management Actions

- Common results across studies
 - Various SRKW behavioral changes occur when vessels are within close proximity (0 m to ≥ 400 m)



Vessel Regulations (2011)





Vessel Presence and Noise




Bioenergetic model to assess the biological significance of responses to vessels


(Noren, Holt, and others)












Lusseau et al. (2009), Noren et al. (2009), Williams et al. (2009)









BEHAVIOR CHANGE

↑ Surface active behaviors
Changes in swim speed
Switching from forage to travel (↑ swim speed)
Vocal modifications

Change in metabolic rate

LIFE FUNCTION
IMMEDIATELY
AFFECTED
Prey intake

-Increased DPERs







Noren et al. (2009), Williams et al. (2009)



Energetic Cost of Surface Active Behaviors








SABs can increase energy expenditure


- Breaches are energetically costly
- Tail slaps are less costly




Noren, Dunkin, Williams (unpublished data)



Energetic Cost of Increasing Swim Speed





* From Noren (2011)

| | FMR at SRKW forage speed (1.1 ms ^{-1*}) | FMR at SRKW travel speed (2.2 ms ^{-1*}) |
|--|---|---|
| Adult Males (4434 kg*) | 1153.7 MJ day ⁻¹ (7.25 X Kleiber BMR) | 1186.1 MJ day ⁻¹ (7.45 X Kleiber BMR) |
| Adult Females (without calves, 3338 kg*) | 636.6 MJ day ⁻¹ (4.95X Kleiber BMR) | 649.7 MJ day ⁻¹ (5.05X Kleiber BMR) |

Williams and Noren (2009)




Do Behavioral Responses Significantly Increase Daily Prey Energy Requirements (DPERs)?













Conclusions

- Research has directly influenced management actions
 - Vessel regulations (2011)
- Current work assesses energetic impacts of vessel presence
 - Are SRKW DPERs (Noren 2011) significantly greater in the presence of vessels?
- Future work could assess the efficacy of vessel regulations in reducing SRKW behavioral responses
 - Too costly to repeat behavioral studies, given budget restrictions
 - Data collected during DTAG studies might provide insight into changes in activity states, but sample size may be limited









Relevant Peer-Reviewed Publications Funded by NOAA NWFSC

Southern Resident Killer Whale Behavioral Responses to Vessels

- *Lusseau D., Bain D.E., Williams R., Smith J.C. (2009) Vessel traffic disrupts the foraging behavior of southern resident killer whales *Orcinus orca*. Endangered Species Research 6:211–221.
- *Noren D.P., Johnson A.H., Rehder D., Larson A. (2009) Close approaches by vessels elicit surface active behaviors by Southern Resident killer whales. Endangered Species Research 8:179-192.
- *Williams R., Bain D.E., Smith J.C., Lusseau D. (2009) Effects of vessels on behaviour patterns of individual southern resident killer whales *Orcinus orca*. Endangered Species Research 6:199-209.

Southern Resident Killer Whale Summer Critical Habitat Use and Potential Marine Protected Areas

Ashe E., Noren D.P., Williams R. (2010) Animal behaviour and marine protected areas: Incorporating behavioural data into the selection of marine protected areas for an endangered killer whale population. Animal Conservation 13:196-203.

Noren D.P. and Hauser D.D.W. (in review) Surface-based observations can assess activity budgets and fine-scale habitat use by an endangered killer whale population.

Northern Resident Killer Whale Behavioral Responses to Vessels

Williams R. and Ashe E. (2007) Killer whale evasive tactics vary with boat number. Journal of Zoology 272:390-397.

Resident Killer Whale Energetics

Noren D.P. (2011) Estimated field metabolic rates and prey requirements of resident killer whales. Marine Mammal Science 27:60–77.

Williams R. and Noren D.P. (2009) Swimming speed, respiration rate and estimated cost of transport in adult killer \whales. Marine Mammal Science 25: 327–350.

*Denotes open access

